

## 2011 REVISED ROUTINE PUBLICLY FUNDED VACCINE SCHEDULE FOR HEALTHY CHILDREN

AGE	VACCINE # 1	VACCINE # 2	VACCINE #3
2 months	Pediacel # 1	Prevnar®13 #1 (Low Risk)	Rotarix #1 (oral)
4 months	Pediacel #2	Prevnar®13 #2 (Low Risk)	Rotarix #2 (oral)
6 months	Pediacel #3		
12 months	MMR	Menjugate	Prevnar®13#3 (Low Risk)
15 months	Varicella		
18 months	Pediacel #4		
4 years	Adacel®Polio	MMRV (Priorix-Tetra)	

**Rotarix** = Recommended to prevent gastroenteritis caused by a rotavirus infection in young children. Can start after 6 weeks of age and completed by 24 weeks of age.

**Pediacel** = diphtheria, pertussis, polio, tetanus, Hib

**MMR** = measles, mumps, rubella (german measles) First dose **on or after 1<sup>st</sup> birthday**.

**Priorix-Tetra (MMRV)** = measles, mumps, rubella (german measles) and varicella

**Adacel®Polio** = diphtheria, pertussis, polio, tetanus

**Menjugate (Meningococcal C)** = One dose if > 1 year. One-year-old children immunized in infancy (i.e. < 1 year of age) require another dose at least one year after the last dose for enhanced protection.

**Prevnar®13** = A three dose schedule for healthy (low risk) children - 2 primary doses at 2 and 4 months of age and a booster dose at 12 months. For high risk children the schedule is three doses at two, four and six months and then a booster at 15 months. Refer to Prevnar 13 Eligibility Schedule (Tables 2 and 4)

**Quadracel** = **Only for 5 or 6 year old unimmunized children starting their primary series.**